

HUMAN TNFSF12 PROTEIN, HFC TAG

货号: 11223

产品全名: 人 TNFSF12 蛋白

规格: 10/50/100 µg

基因符号 APO3L;DR3LG;TNLG4A;TWEAK

目标蛋白: TNFSF12

UNIPROT ID: O43508

描述: Recombinant Human TNFSF12 Protein with N-terminal Human Fc tag

背景: The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein is a ligand for the FN14/TWEAKR receptor. This cytokine has overlapping signaling functions with TNF, but displays a much wider tissue distribution. This cytokine, which exists in both membrane-bound and secreted forms, can induce apoptosis via multiple pathways of cell death in a cell type-specific manner. This cytokine is also found to promote proliferation and migration of endothelial cells, and thus acts as a regulator of angiogenesis. Alternative splicing results in multiple transcript variants. Some transcripts skip the last exon of this gene and continue into the second exon of the neighboring TNFSF13 gene, such read-through transcripts are contained in GeneID 407977, TNFSF12-TNFSF13.

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 43.3 kDa after removal of the signal peptide.

分子特征: hFc(Glu99-Ala330) TNFSF12(Ser94-His249)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human TNFSF12 Protein, hFc Tag on SDS-PAGE under reducing condition.

Human TNFSF12, hFc tagged protein ELISA
0.1 μ g of Human TNFSF12, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 1 μ g/ml (100 μ l/well) Human TNFSF12 Protein, hFc Tag (11223) can bind Anti-TNFSF12 Rabbit mAb in a linear range of 3.2-80 ng/mL.