

IDH1(R132P)

产品名称: IDH1(R132P)

货号: 26406

基因符号: IDH2; D2HGA2; ICD-M; IDH; IDHM; IDP; IDPM; mNADP-IDH

描述: IDH1(R132P) 小鼠单抗

背景: Isocitrate dehydrogenases (IDH) catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. Mutations affecting Arg-132 are tissue-specific, and suggest that this residue plays a unique role in the development of high-grade gliomas. Mutations of Arg-132 to Cys, His, Leu or Ser abolish magnesium binding and abolish the conversion of isocitrate to alpha-ketoglutarate. Instead, alpha-ketoglutarate is converted to R-2-hydroxyglutarate. Elevated levels of R-2-hydroxyglutarate are correlated with an elevated risk of malignant brain tumors.

免疫原: A synthetic peptide from the internal region of IDH1 which includes the mutation of R132P, human origin.

经过测试的应用: ELISA, WB, IF, IHC

推荐稀释比: ELISA: 1:1000-1:5000WB: 1:100-1:1000IF: 1:50-1:100IHC: 1:50-1:100

Concentration: 1 mg/ml

种属反应性: Mouse

形式: Liquid

克隆性: Monoclonal

亚型: IgG


纯化: Purified from ascites

Preservative: No

成分: PBS (without Mg^{2+} and Ca^{2+}), pH 7.4, 150 mM NaCl, 50% glycerol

种属反应性: recognizes R132P mutant, but not wild type IDH2 of vertebrates.

储存条件: Store at $-20^{\circ}C$. Avoid repeated freezing and thawing



蛋白质印迹: WB: anti-IDH1 (R132P) mAb **Immunofluorescence of cells expressing IDH1 proteins with Anti-IDH1(R132P) antibody.** HEK293T cells were transfected with pCDNA3-GFP-IDH1(R132P) plasmid, pCDNA3-GFP-IDH1 (WT) plasmid, pCDNA3-GFP-IDH1(R132S) plasmid, pCDNA3-GFP-IDH1(R132L) plasmid, pCDNA3-GFP-IDH1(R132G) plasmid or pCDNA3-GFP-IDH1(R132C) plasmid, then fixed and stained with Anti-IDH1(R132P) monoclonal antibody (货号:26406).