

## IDH2(R172G)

产品名称: IDH2(R172G) 小鼠单抗

货号: 26231

基因符号: IDH2; D2HGA2; ICD-M; IDH; IDHM; IDP; IDPM; mNADP-IDH

描述: IDH2(R172G) 小鼠单抗

背景: Isocitrate dehydrogenase (IDH) catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. The isocitrate and isopropylmalate dehydrogenases family has three members, IDH1, IDH2 and IDH3. IDH2 plays a role in intermediary metabolism and energy production. Defects in IDH2 are the cause of D-2-hydroxyglutaric aciduria type 2 (D2HGA2). Somatic mosaic mutations of this protein have also been found associated to Ollier disease and Maffucci syndrome, and R172G IDH2 mutations do exist in diffusely infiltrative gliomas.

免疫原: A synthetic peptide from the internal region of IDH2 which includes the mutation of R172G, human origin.

经过测试的应用: ELISA, WB, IF, IHC

推荐稀释比:

ELISA: 1:1000-1:5000

WB: 1:100-1:1000

IF: 1:50-1:100

IHC: 1:50-1:100

**Concentration:** 1 mg/ml

种属反应性: **Mouse**

形式: Liquid

克隆性: Monoclonal

亚型: **IgG**

纯化: Purified from ascites

成分: PBS (without  $Mg^{2+}$  and  $Ca^{2+}$ ), pH 7.4, 150 mM NaCl, 50% glycerol


种属反应性: recognizes R172G mutant, but not wild type IDH2 of vertebrates.

储存条件: Store at  $-20^{\circ}C$ . Avoid repeated freezing and thawing

蛋白质印迹:

## IDH2 proteins

R172G WT




WB: Anti-IDH2(R172G) mAb

### Western blot analysis of recombinant IDH2(R172G) and wild type proteins.

Purified His-tagged IDH2(R172G) (lane 1) and corresponding wild type IDH2 protein (lane 2) were blotted with Anti-IDH2(R172G) monoclonal antibody (货号:26231).

### Immunofluorescence:



### Immunofluorescence of cells expressing IDH2 proteins with Anti-IDH2(R172G) antibody.

HEK293T cells were transfected with pCDNA3-GFP-IDH2 (WT) plasmid (left column) or pCDNA3-GFP-IDH2(R172G) plasmid (right column), then fixed and stained with Anti-IDH2(R172G) monoclonal antibody (货号:26231).