

IL5RA (DMC393) IGG1 CHIMERIC MAB

货号: 28211

产品全名: IL5RA(DMC393) IgG1 Chimeric 单抗

基因符号 IL5Ra;CD125;IL-5 R alpha

描述: IL5RA antibody(DMC393) IgG1 Chimeric 单抗

背景: The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3); colony stimulating factor 2 (CSF2:GM-CSF); and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding; and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin); which is required for IL5 mediated activation of the transcription factor SOX4. Several alternatively spliced transcript variants encoding four distinct isoforms have been reported.

经过测试的应用: Flow Cyt

推荐稀释比: Flow Cyt 1:100

种属反应性: Rabbit

亚型: Rabbit:Human Fc chimeric IgG1

纯化: Purified from cell culture supernatant by affinity chromatography

种属反应性: 人 IL5RA

成分: Lyophilized from sterile PBS, pH 7.4. 5 % – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form.

After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).




Figure 1. Flow cytometry analysis with Anti-IL5RA (DMC393) on Expi293 cells transfected with human IL5RA (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).