

MOUSE CD47 PROTEIN, HFC TAG

货号: 12136

产品全名: Mouse CD47 蛋白

规格: 10/50/100 µg

基因符

号 9130415E20Rik;AA407862;A1848868;AW108519;B430305P08Rik;IAP;Itgp

目标蛋白: CD47

UNIPROT ID: Q61735

描述: Recombinant mouse CD47 protein with C-terminal human Fc tag

背景: Has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. Plays an important role in memory formation and synaptic plasticity in the hippocampus. Receptor for SIRPA, binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Interaction with SIRPG mediates cell-cell adhesion, enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation. May play a role in membrane transport and/or integrin dependent signal transduction. May prevent premature elimination of red blood cells. May be involved in membrane permeability changes induced following virus infection (By similarity).[UniProtKB/Swiss-Prot Function]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 40.0 kDa after removal of the signal peptide. The apparent molecular mass of mCD47-hFc is approximately 55-70 kDa due to glycosylation.

分子特征: Mouse CD47(Gln19-Lys140) hFc(Glu99-Ala330)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Mouse CD47 Protein, hFc Tag on SDS-PAGE under reducing condition.