

## RAB7(Q67L) PROTEIN

产品名称: Rab7(Q67L) 突变蛋白

货号: 10135

产品全名: Rab7 Protein Q67L 突变蛋白

基因符号 Member RAS oncogene family, Rab7a, MGC102153

**Source:** Human, recombinant full length, His6-tag

**Expression 种属反应性:** E. coli

分子量: 23 kDa

纯化: >95% by SDS-PAGE

**Introduction:** Members of the Rab family of Ras-related GTP-binding proteins are important regulators of vesicular transport and are located in specific intracellular compartments. Rab7 has been localized to late endosomes and shown to be important in the late endocytic pathway. In addition, it has been shown to have a fundamental role in the cellular vacuolation induced by the cytotoxin VacA of *Helicobacter pylori*.

**Amino Acid Sequence** (1-207, Q67L)

**MTSRKKVLLKVIILGDSGVGKTSLMNQYVNKKFSNQYKATIGADFLTKEVMVDDRLVTMQIWDTAGL  
ERFQSLGVAFYRGADCCVLVFDVTAPNTFKTLDSWRDEFLIQASPRDPENFPFVVLGNKIDLENRQV  
ATKRAQAWCYSKNNIPYFETSAKEAINVEQAFQTIARNALKQETEVELYNEFPEPIKLDKNDRAKASAE  
CSC**

### Properties

**Physical Appearance (form):** Dissolved in 20mM Tris-HCl, pH8.0, 150mM NaCl.


**Physical Appearance (form):** White or clear

**Concentration:** 1 mg/mL

**Storage:** -80°C

### Preparation Instructions:

Centrifuge the vial before open the cap and reconstitute in water. Adding of 10 mM  $\beta$ -mercaptoethanol or 1 mM DTT into the solution to protect the protein is recommended and using of non-ionic detergents such as n-Dodecyl  $\beta$ -D-maltoside (DoDM) or polyethylene detergents (e.g. C12E10) also help to stabilize the protein. Avoid repeated freezing and thawing after reconstitution. The purity of His-tagged Rab7 Q67L was determined by SDS-PAGE and Coomassie Brilliant Blue Staining.



**References:**

1. Edinger, A. L. et al., Dev. Cell 5: 571-582, 2003.
2. Houlden, H. et al., Ann. Neurol. 56: 586-590, 2004.
3. Meggouh, F. et al., Neurology 67: 1476-1478, 2006.
4. Rak, A. et al., Cell 117: 749-760, 2004.
5. Verhoeven, K. et al., Am. J. Hum. Genet. 72: 722-727, 2003.