

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## SARS-COV-2 (2019-NCOV) S PROTEIN RBD, MFC TAG

货号: 11301

产品全名: SARS-CoV-2 (2019-NCoV) S Protein RBD

规格: 10/50/100 µg

基因符号 S protein RBD;Spike glycoprotein Receptor-binding domain;S glycoprotein RBD;Spike

protein RBD;COVID-19 目标蛋白: S protein RBD **UNIPROT ID:** P0DTC2

描述: Recombinant SARS-CoV-2 (2019-nCoV) S protein RBD with C-terminal mouse Fc tag 背景: SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which accounts for recognizing the cell surface receptor, ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell

物种/宿主: HEK293

response.

分子量: The protein has a predicted molecular mass of 51.3 kDa after removal of the signal peptide.

分子特征: S protein RBD(Arg319-Phe541) mFc(Pro99-Lys330)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. 储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010




Figure 1. SARS-CoV-2 (2019-nCoV) S protein RBD, mFc Tag on SDS-PAGE under reducing condition.

## S-RBD, mFc Tagged protein ELISA

0.2 µg of S-RBD, mFc Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) S-RBD, mFc tagged protein (11301) can bind Human ACE2, hFc Tagged protein 11196 in a linear range of 7.81-87.7 ng/ml.