

SARS-COV-2 RBD (DM35) RABBIT MAB

货号: 28324

产品全名: SARS-CoV-2 RBD(DM35) 兔单克隆抗体

基因符号 SARS-CoV-2 RBD

描述: SARS-CoV-2 RBD antibody(DM35) 兔单克隆抗体

背景: SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits; S1 and S2. S1 mainly contains a receptor binding domain (RBD); which accounts for recognizing the cell surface receptor; ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response.

经过测试的应用: ELISA; Flow Cyt

推荐稀释比: ELISA 1:5000-10000; Flow Cyt 1:100

种属反应性: Rabbit

亚型: Rabbit IgG

纯化: Purified from cell culture supernatant by affinity chromatography

种属反应性: SARS-CoV-2 S protein RBD

成分: Lyophilized from sterile PBS, pH 7.4. 5% - 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).




Figure 1. Elisa plate pre-coated by 2 µg/ml (100µl/well) SARS-CoV-2 RBD protein can bind Rabbit Anti-SARS-CoV-2 RBD monoclonal antibody (clone:DM35) in a linear range of 0.19-200 ng/ml.




Figure 2. Competition flow cytometry assay demonstrating Rabbit anti-RBD monoclonal antibody (clone: DM35) blockade of SARS-CoV-2 (COVID-19) S protein RBD (1µg/ml, [getskuurl sku=11301 binding to Expi 293 cell line transfected with human ACE2. IC50=1013ng/ml. The Y-axis represents the geometric mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.