

HUMAN SLC1A5 FULL LENGTH PROTEIN

货号: 11038

产品全名: 人 SLC1A5 Full Length 蛋白

规格: 10/50/100 µg

基因符号 AAAT; ASCT2; ATBO; M7VI; M7VSI; R16; RDRC

目标蛋白: SLC1A5

UNIPROT ID: Q15758

描述: 人 SLC1A5 full length protein-synthetic nanodisc

背景: The SLC1A5 gene encodes a sodium-dependent neutral amino acid transporter that can act as a receptor for RD114/type D retrovirus (Larriba et al., 2001 [PubMed 11781704]).[supplied by OMIM, Jan 2011]

物种/宿主: HEK293

分子量: The human full length SLC1A5 protein has a MW of 56.4 kDa

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

ELISA assay to evaluate SLC1A5-Nanodisc
0.2µg Human SLC1A5-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag SLC1A5-Nanodisc (0.2 µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with SLC1A5-Nanodisc is 3.850ng/ml.



Figure2. Human SLC1A5-Nanodisc, Flag Tag on SDS-PAGE